

AD-A077 888

NORTH CAROLINA UNIV AT CHAPEL HILL INST OF STATISTICS F/G 12/1  
ON THE ASYMPTOTIC NORMALITY OF ROBUST REGRESSION ESTIMATES.(U)  
JUL 79 R J CARROLL , D RUPPERT NSF-MCS78-01240

UNCLASSIFIED

MIMEO SER-1242

AFOSR-TR-79-1101

NL

1 OF 1  
ADA  
077888




END  
DATE  
FILMED

1 -80


DDC





② LEVEL II

AD A 077888



*See 1472*

On the Asymptotic Normality of Robust Regression Estimates

by

Raymond J. Carroll  
and  
David Ruppert

University of North Carolina at Chapel Hill

Institute of Statistics Mimeo Series #1242

July 1979

79 11 27 058

DEPARTMENT OF STATISTICS  
Chapel Hill, North Carolina

DDC  
RECEIVED  
DEC 11 1979  
B


DDC FILE COPY

Approved for public release;  
distribution unlimited.

ON THE ASYMPTOTIC NORMALITY OF ROBUST REGRESSION ESTIMATES

by

Raymond J. Carroll\* and David Ruppert\*\*  
University of North Carolina at Chapel Hill



Abstract

Huber's (1973) proof of asymptotic normality of robust regression estimates is modified to include the estimates used in practice, which have unknown scale and only piecewise smooth defining functions  $\psi$ .

Key Words and Phrases: Robustness, M-estimates, regression, linear model, asymptotic theory.

\*This research was supported by the Air Force Scientific Office of Scientific Research under contract AFOSR-75-2796.

\*\*This research was supported by the National Science Foundation Grant NSF MCS78-01240.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer



## 1. Introduction

We consider the general linear model

$$(1.1) \quad y_i = \tau_i + z_i/\sigma_0 \quad (i = 1, \dots, n),$$

where

$$(1.2) \quad \tau_i = \sum_{j=1}^p c_{ij} \beta_j^{(0)}.$$

Here the  $\beta_j^{(0)}$  are unknown parameters, the  $c_{ij}$  are known constants, the  $z_i$  are independent and identically distributed with common distribution function  $F$  (symmetric about zero), and  $\sigma_0$  is a constant to be chosen later. Huber (1973) proposed estimating  $\beta$  by solving the system of equations

$$(1.3) \quad \sum_i \psi(\sigma(y_i - t_i(\beta))) c_{ij} = 0 \quad (j = 1, \dots, p)$$

$$(1.4) \quad \sum_i \{\psi^2(\sigma(y_i - t_i(\beta))) - \xi\} = 0,$$

where

$$t_i(\beta) = \sum_{j=1}^p c_{ij} \beta_j$$

$$\xi = E_{\phi} \psi^2(z),$$

(the expectation being taken under the standard normal distribution) and  $\psi$  is a monotone nondecreasing function. Let  $C = \{c_{ij}\}$ ,  $\Gamma = C(C^T C)^{-1} C^T$  be the projection matrix, and  $\epsilon$  be the maximum diagonal element of  $\Gamma$ .

Huber (1973) considers  $\sigma$  fixed and known ( $\sigma \equiv 1$ ) so that one only need solve (1.3). He proves that if  $\epsilon p^2 \rightarrow 0$  as  $n \rightarrow \infty$  (which implies  $p^3/n \rightarrow 0$ ) and if  $\psi$  is bounded with two continuous bounded derivatives, then all estimates of the form  $\hat{\alpha} = \sum_{j=1}^p a_j \hat{\beta}_j$  ( $\sum_j a_j^2 = 1$ ) are asymptotically normal. It is

routine to extend his results to the system (1.3), (1.4) (see below). The smoothness conditions on  $\psi$  are not satisfied for three of the most commonly used functions, namely

Huber's function

$$\begin{aligned}\psi(x) &= x & |x| < c \\ &= c \operatorname{sign}(x) & |x| \geq c\end{aligned}$$

Hampel's function

$$\begin{aligned}\psi(x) &= -\psi(-x) = x & 0 < x < a \\ &= a & a \leq x < b \\ &= a \left( \frac{c-x}{c-b} \right) & b \leq x < c \\ &= 0 & x \geq c\end{aligned}$$

Andrew's function

$$\begin{aligned}\psi(x) &= -\psi(-x) = \operatorname{sine}(x/c) & 0 < x < \pi c \\ &= 0 & x \geq \pi c.\end{aligned}$$

In this note we weaken Huber's conditions to include the common  $\psi$ -functions, at the cost of slightly strengthened conditions on the rate of growth of  $p$ , the dimension of the problem. The results have been applied by Carroll and Ruppert (1979) to the problem of testing for heteroscedasticity (Bickel (1978)).

As in Huber's proof, we assume  $C'C = I$ . Because of the invariance of the problem, we can take  $\beta_j^{(0)} = 0$  ( $j = 1, \dots, p$ ) and  $\sigma_0 = 1$ , primarily to simplify notation.

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
ODC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION _____	
BY _____	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

## 2. Notation, Assumptions and Main Results

Let  $a$  be an arbitrary  $(p \times 1)$  vector for which  $\sum a_j^2 = \|a\|^2 = 1$ . Define  $s_i = \sum_{j=1}^p c_{ij} a_j$ . Note that  $\|t(\beta)\|^2 = \|\beta\|^2$ . Define for  $j = 1, \dots, p$ ,

$$\phi_j(\beta, \sigma) = - \sum_{i=1}^n \psi(\sigma(y_i - t_i(\beta))) c_{ij} / E\psi'(y_1)$$

$$\psi_j(\beta, \sigma) = \beta_j - \sum_{i=1}^n \psi(y_i) c_{ij} / E\psi'(y_1),$$

while

$$\phi_{p+1}(\beta, \sigma) = -n^{-1/2} \sum_{i=1}^n \{\psi^2(\sigma(y_i - t_i(\beta))) - \xi\} / 2 E y_1 \psi(y_1) \psi'(y_1)$$

$$\psi_{p+1}(\beta, \sigma) = -n^{-1/2} \left( (\sigma-1) + \sum_{i=1}^n \{\psi^2(y_i) - \xi\} / 2 E y_1 \psi(y_1) \psi'(y_1) \right).$$

Our estimates are solutions to  $\phi_j(\hat{\beta}, \hat{\sigma}) = 0$  ( $j = 1, \dots, p+1$ ), and we hope to approximate  $(\hat{\beta}, \hat{\sigma})$  by  $(\tilde{\beta}, \tilde{\sigma})$  which solves  $\psi_j(\tilde{\beta}, \tilde{\sigma}) = 0$  ( $j = 1, \dots, p+1$ ).

We make the following assumptions.

(2.1)  $\psi$  is odd, bounded, and constant outside a finite interval.

(2.2)  $\psi$  is Lipschitz of order one and has two continuous bounded derivatives except at a finite number of points, which we take without loss as  $\pm c$ .

(2.3)  $F$  is symmetric about zero.

(2.4)  $F$  is Lipschitz in neighborhoods of  $\pm c$ .

(2.5)  $E \psi'(y_1) \neq 0$ ,  $E y_1 \psi(y_1) \psi'(y_1) \neq 0$ .

Theorem 1. If (2.1) - (2.5) hold and, in addition, there is a sequence  $a_n \rightarrow 0$  such that



$$(2.6) \quad (\epsilon p / a_n^2) \rightarrow 0, \quad (\epsilon n a_n) \rightarrow 0,$$

then there is a sequence of solutions to (1.3) - (1.4) such that

$$(2.7) \quad ||(\hat{\beta}, (np)^{1/2} (\hat{\sigma} - 1))|| = o_p(p).$$

If in addition

$$(2.8) \quad (\epsilon p^2 / a_n^2) \rightarrow 0, \quad (\epsilon n p a_n) \rightarrow 0,$$

then

$$(2.9) \quad ||(\hat{\beta}, (np)^{1/2} (\hat{\sigma} - 1)) - (\tilde{\beta}, (np)^{1/2} (\hat{\sigma} - 1))|| \xrightarrow{P} 0.$$

Thus, (2.1) - (2.5) and (2.8) imply that all estimates of the form

$$\hat{\alpha} = \sum_{j=1}^p \alpha_j \hat{\beta}_j \quad (||\alpha||^2 = 1) \text{ are asymptotically normal.}$$

Remark: The result (2.7) is the starting point in constructing robust tests for heteroscedasticity (Bickel (1978), Carroll and Ruppert (1979)); it implies the crucial assumption T of Bickel's Theorem 3.1. For balanced designs ( $\epsilon = p/n$ ), assumption (2.6) is satisfied by choosing  $a_n = p^{-(1+\gamma)}$  for small  $\gamma > 0$ , which then requires  $p^{4+2\gamma}/n \rightarrow 0$ , as compared to Huber's condition  $p^2/n \rightarrow 0$  when  $\psi$  has two derivatives.

### 3. Proofs

Proposition 1. Suppose that (2.1) - (2.5) hold, that  $\psi$  has two bounded continuous derivatives, and that  $\epsilon p \rightarrow 0$ . Then on the set

$$||(\beta, (pn)^{1/2} (\sigma - 1))||^2 \leq Kp,$$

the following hold:



$$(3.1) \quad ||\Phi(\beta, \sigma) - \Psi(\beta, \sigma)|| = O_p((\epsilon p^2)^{1/2}),$$

$$(3.2) \quad ||\Phi(\beta, \sigma) - (\beta, (np)^{1/2}(\sigma - 1))|| = O_p(p^{1/2} + (\epsilon p^2)^{1/2}).$$

Proof of Proposition 1. By a Taylor series expansion,

$$(3.3) \quad \left| \sum_{j=1}^p a_j (\Phi_j(\beta, \sigma) - \Psi_j(\beta, \sigma)) + (\sigma - 1) \sum_{i=1}^n s_i y_i \psi'(y_i) / E \psi'(y_1) - \sigma \sum_{i=1}^n s_i t_i(\beta) (\psi'(y_i) - E \psi'(y_1)) / E \psi'(y_1) \right|$$

$$= |A_n|$$

$$\leq |\sigma - 1| \left| \sum_{i=1}^n s_i t_i(\beta) y_i \psi''((1 + \eta_{2i}) y_i) / E \psi'(y_1) \right|$$

$$+ \sigma^2 \left| \sum_{i=1}^n s_i t_i^2(\beta) \psi''(\sigma y_i + \eta_{1i}) / E \psi'(y_1) \right|.$$

On the set  $||\beta||^2 \leq Kp$ ,  $|\sigma - 1| \leq 1/2$ ,  $||a|| = 1$ , Huber shows that the second term on the r.h.s. of (3.3) is  $O(\epsilon^{1/2} ||\beta||^2)$ . Since  $\psi'' = 0$  outside a finite set and  $|\eta_{2i}| > 1/4$ , the first term on the r.h.s. of (3.3) is bounded by

$$M |\sigma - 1| (\sum s_i^2)^{1/2} (\sum t_i^2(\beta))^{1/2} = O(||\beta|| |\sigma - 1|).$$

Thus,

$$|A_n| = O(\epsilon^{1/2} ||\beta||^2 + |\sigma - 1| ||\beta||).$$

We next consider  $A_n$ , in particular its last two terms. Since  $\sum s_i^2 = 1$  and  $E y_1 \psi'(y_1) = 0$ ,

$$(3.4) \quad (\sigma - 1) \sum_{i=1}^n s_i y_i \psi'(y_i) / E \psi'(y_1) = O_p(|\sigma - 1|).$$

Huber shows that

$$(3.5) \quad \sum_{i=1}^n s_i t_i(\beta) (\psi'(y_i) - E \psi'(y_1)) / E \psi'(y_1) = O_p(\epsilon^{1/2} ||\beta||^2).$$

Thus, uniformly on the set  $||\beta||^2 \leq Kp$ ,  $|\sigma - 1| \leq Kn^{-1/2}$ ,  $||a|| = 1$  we have from (3.3) - (3.5)

$$(3.6) \quad \left| \sum_{j=1}^p a_j (\phi_j(\beta, \sigma) - \psi_j(\beta, \sigma)) \right| = O_p(\epsilon^{1/2} p + pn^{-1/2}).$$

Another Taylor series expansion shows that

$$(3.7) \quad |B_n| = |\phi_{p+1}(\beta, \sigma) - \psi_{p+1}(\beta, \sigma) + C_{n1} + C_{n2}| \\ \leq M\{|\sigma - 1| ||\beta|| + pn^{-1/2} + n^{-1/2} |\sigma - 1|^2\},$$

where

$$-\{2 E y_1 \psi(y_1) \psi'(y_1)\} C_{n1} = 2n^{-1/2} (\sigma - 1) \sum_{i=1}^n (y_i \psi(y_i) \psi'(y_i) - E y_1 \psi(y_1) \psi'(y_1))$$

and

$$\{2 E y_1 \psi(y_1) \psi'(y_1)\} C_{n2} = 2 \sigma n^{-1/2} \sum_{i=1}^n t_i(\beta) \psi(y_i) \psi'(y_i).$$

Since  $C_{n1} = O_p(|\sigma - 1|)$  and  $C_{n2} = O_p((p/n)^{1/2})$ , we obtain

$$|\phi_{p+1}(\beta, \sigma) - \psi_{p+1}(\beta, \sigma)| = O_p(pn^{-1/2}).$$

Since  $(\epsilon p^2)^{1/2} \geq (p^3/n)^{1/2} \geq pn^{-1/2}$ , we thus obtain (3.1). Equation (3.2) follows from (3.20) of Huber (1973).

Proof of Theorem 1. The proof of Proposition 1 makes it clear that we need to obtain bounds for  $|A_n|$  and  $|B_n|$  uniformly on the set  $||\beta|| \leq Kp$ ,  $|\sigma - 1| \leq Kn^{-1/2}$  and  $||a|| = 1$ . Rewrite

$$E \psi'(y_1) |A_n| = \left| \sum_{i=1}^n H(i, \beta, \sigma) \left\{ \begin{aligned} & s_i (\psi(\sigma(y_i - t_i(\beta))) - \psi(y_i)) \\ & + (\sigma - 1) s_i y_i \psi'(y_i) \\ & - \sigma s_i t_i(\beta) (\psi'(y_i) - E \psi'(y_1)) \end{aligned} \right\} \right|.$$

Let  $I$  be the indicator function and let  $a_n \rightarrow 0$ ,  $n^{1/2} a_n \rightarrow \infty$ . Then

$$\begin{aligned} |A_n| &= \left| \sum_{i=1}^n H(i, \beta, \sigma) \left\{ \begin{aligned} & I(-c + a_n \leq y_i \leq c - a_n) \\ & + I(y_i \geq c + a_n) + I(y_i \leq -c - a_n) \\ & + I(-c - a_n < y_i < -c + a_n) + I(c - a_n < y_i < c + a_n) \end{aligned} \right\} \right| \\ &= |A_{n1} + A_{n2} + A_{n3} + A_{n4} + A_{n5}|. \end{aligned}$$

For notational purposes, define  $d_i = \sigma(y_i - t_i(\beta))$ . Then

$$\begin{aligned} A_{n1} &= \sum_{i=1}^n H(i, \beta, \sigma) I(-c + a_n < y_i < c - a_n) \\ &\times \{I(-c < d_i < c) + I(|d_i| > c)\} = A_{n1}^{(1)} + A_{n1}^{(2)}. \end{aligned}$$

By Proposition 1,  $|A_{n1}^{(1)}| = O(\epsilon^{1/2} p)$ . Since  $\psi$  is Lipschitz and constant outside a finite interval,

$$|A_{n1}^{(2)}| \leq M \sum_{i=1}^n |s_i| \{|\sigma - 1| + |t_i(\beta)|\} I(-c + a_n < y_i < c - a_n \mid |d_i| > c).$$

However, since  $|\sigma - 1| \leq K n^{-1/2}$  and  $n^{1/2} a_n \rightarrow 0$ ,

$$|A_{n1}^{(2)}| \leq M \epsilon^{1/2} \sum_{i=1}^n \{|\sigma - 1| + |t_i(\beta)|\} I\{|t_i(\beta)| > a_n/2\}.$$

Now

$$(3.8) \quad \sum_{i=1}^n I\{|t_i(\beta)| > a_n/2\} \leq 4p/a_n^2$$



$$(3.9) \quad \sum_{i=1}^n |t_i(\beta)| I\{|t_i(\beta)| > a_n/2\} \leq 4p/a_n,$$

so that (3.8) and (3.9) imply

$$|\Lambda_{n1}^{(2)}| \leq 4M \epsilon^{1/2} \{|\sigma - 1| p/a_n^2 + p/a_n\}.$$

However,  $|\sigma - 1|/a_n \leq K(na_n^2)^{-1/2}$  so that

$$(3.10) \quad |\Lambda_{n1}| = O(\epsilon^{1/2} p/a_n (1 + (na_n^2)^{-1/2})).$$

The same bound (3.10) holds for  $|\Lambda_{n2}|$  and  $|\Lambda_{n3}|$ . Noting once again that  $\psi$  is Lipschitz and constant outside a finite interval,

$$\begin{aligned} |\Lambda_{n5}| &\leq \sum_{i=1}^n |s_i| \{|\sigma - 1| + |t_i(\beta)|\} I\{|y_i - c| < a_n\} \\ &\leq M\epsilon^{1/2} \{|\sigma - 1| G_n + p^{1/2} G_n^{1/2}\}, \end{aligned}$$

where

$$(3.11) \quad G_n = \sum_{i=1}^n I\{|y_i - c| < a_n\}.$$

From Lemma 1 of Carroll (1978), provided  $na_n \geq \log n$ ,

$$G_n = O_p(na_n).$$

This gives

$$(3.12) \quad |\Lambda_{n5}| = O_p((\epsilon npa_n)^{1/2}).$$

The bound (3.12) also holds for  $|\Lambda_{n4}|$ . Thus,

$$(3.13) \quad |\Lambda_n| = O_p(\epsilon^{1/2} p/a_n (1 + (na_n^2)^{-1/2}) + (\epsilon npa_n)^{1/2}).$$

The same bound holds for  $|B_n|$ . Thus,



$$\begin{aligned}
 (3.14) \quad ||\Phi(\beta, \sigma) - \Psi(\beta, \sigma)|| &= O_p(c^{1/2} p(1 + (na_n^2)^{-1/2})/a_n + (c npa_n)^{1/2}) \\
 &= O_p(r(p, n)) .
 \end{aligned}$$

Equation (3.14) is the generalization of Huber's (3.18). As he shows, for sufficiently large  $K$ , on the set  $||\beta||^2 \leq Kp$ ,  $|\sigma - 1| \leq Kn^{-1/2}$ ,

$$(3.15) \quad ||\Phi(\beta, \sigma) - (\beta, (np)^{1/2}(\sigma - 1))|| \leq r(p, n) + \frac{1}{2}(Kp)^{1/2} .$$

Thus, if

$$(3.16) \quad r(p, n)/p^{1/2} \rightarrow 0 ,$$

Brouwer's fixed point theorem enables us to conclude that  $\Phi$  has a zero inside the ball  $||(\beta, (np)^{1/2}(\sigma - 1))|| < Kp$ . Equation (3.16) is true if (2.6) is true. The rest of the proof parallels that of Huber.  $\square$

### References

- Bickel, P.J. (1978). Using residuals robustly I: tests for heteroscedasticity, non-linearity. *Ann. Statist.* 6, 266-291.
- Carroll, R.J. (1978). On almost sure expansions for M-estimates. *Ann. Statist.* 6, 314-318.
- Huber, P.J. (1973). Robust regression: asymptotics, conjectures and Monte-Carlo. *Ann. Statist.* 1, 799-821.
- Huber, P.J. (1977). Robust Statistical Procedures. SIAM, Philadelphia.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR-79-1101</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER <i>(9) Interim Rept.</i>
4. TITLE (and Subtitle) <b>On the Asymptotic Normality of Robust Regression Estimates.</b>	5. TYPE OF REPORT & PERIOD COVERED <b>Interim</b>	
7. AUTHOR(s) <b>Raymond J. Carroll and David Ruppert</b>	14. PERFORMING ORG. REPORT NUMBER <b>Mimeo Series No. 1242</b>	8. PERFORMING ORG. REPORT NUMBER <b>AFOSR-75-2796</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>University of North Carolina Department of Statistics Chapel Hill, N.C. 27514</b>	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 2304 A5</b>	
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research Bolling AFB, DC 20332</b>	12. REPORT DATE <b>Jul 1979</b>	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>(12) 13</b>	13. NUMBER OF PAGES <b>11</b>	
	15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) <b>Approved for public release - distribution unlimited.</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>Robustness, M-estimates, regression, linear model, asymptotic theory</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>Huber's (1973) proof of asymptotic normality of robust regression estimates is modified to include the estimates used in practice, which have unknown scale and only piecewise smooth defining functions <math>\psi</math>.</b> <i>Lpsi</i>		

DD FORM 1473  
1 JAN 73

EDITION OF NOV 65 IS OBSOLETE

UNCLASSIFIED

410 064  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)